

AD-A049 526

UNIVERSITY OF SOUTHERN CALIFORNIA LOS ANGELES MODELL--ETC F/6 12/1
DIRECT TRANSFORMATION OF VARIATIONAL PROBLEMS INTO CAUCHY SYSTE--ETC(U)
1977 J HESS, R KALABA AFOSR-77-3383

UNCLASSIFIED

USCMR6-7710

AFOSR-TR-78-0016


NL

| OF |
AD
A049526



END
DATE
FILMED
2-78
DDC





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 049526

AFOSR-TR- 78 - 0016




Jr (2)

*see book
page 143*

AD No.

DDC FILE COPY



modelling research group



department of economics
university of southern california
university park
los angeles, california
90007

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)

NOTICE OF TRANSMITTAL TO DDC

This technical report has been reviewed and is approved for public release IAW AFR 190-12 (7b). Distribution is unlimited.

A. D. BLOSE

Technical Information Officer




Working Paper: Not to be quoted without authors' permission.

Author: Linda A. Anderson

DISTRIBUTION STATEMENT A


Approved for public release;
Distribution Unlimited



DIRECT TRANSFORMATION OF VARIATIONAL PROBLEMS
INTO CAUCHY SYSTEMS. I. SCALAR-QUADRATIC CASE*

by

James Hess¹ and Robert Kalaba²




ABSTRACT: This series of papers addresses three interrelated problems: the solution of a variational minimization problem, the solution of integral equations, and the solution of an initial valued system of integrodifferential equations. It will be shown that a large class of minimization problems requires the solution of linear Fredholm integral equations. It has also been shown that the solution of a linear Fredholm integral equation is identical to the solution of a Cauchy system. In this paper, we by-pass the Fredholm integral equations and show that the minimization problem directly implies a solution of a Cauchy system. This first paper in the series looks only at quadratic functionals and scalar functions.

KEY WORDS: Variational Problems, Integral Equations, Parametric Imbedding

* Research sponsored by the Air Force Office of Scientific Research, Air Force Systems Command, USAF, under Grant No. AFOSR-77-3383. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.

¹Assistant Professor of Economics, University of Southern California, Los Angeles, California 90007.

²Professor of Biomedical Engineering and Economics, University of Southern California, Los Angeles, California 90007.



White Section <input checked="" type="checkbox"/>	
Buff Section <input type="checkbox"/>	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	and/or SPECIAL
A	

1. INTRODUCTION

Many interesting optimization problems result in variational problems of finding a function $z(t)$, $0 \leq t \leq 1$, that minimizes the functional

$$W[z] = \lambda \int_0^1 \int_0^1 z(s)k(t,s)z(t)dt ds + \int_0^1 z(t)^2 dt + 2 \int_0^1 f(t)z(t)dt.$$

By standard variational techniques it can be shown that the optimal function, $u(t)$, satisfies a linear Fredholm integral equation (Ref. 1). Recent work in the study of integral equations (Ref. 2) has shown that solutions of integral equations are equivalent to solutions of particular initial valued systems of integrodifferential equations (Cauchy systems). It appears that the three problems -- variational, integral equations, and Cauchy systems -- are equivalent to each other. An important missing link in the analysis has been the demonstration that the variational problem leads directly to a Cauchy system without passing through an integral equation. These papers provide that link. It would seem that the historical emphasis put on the integral equations is but an accident. The Cauchy systems can just as easily be developed and provide interesting insights and computational aid in studying such variational problems.

The direct reduction of the simplest problem in the calculus of variations to a Cauchy system is given in Ref. 3; the train of thought is quite different from that employed here.

2. DERIVATION

Suppose we desire to find a scalar function $z(t)$, $0 \leq t \leq 1$, which minimizes the quadratic functional

$$W[z, \lambda] = \lambda \int_0^1 \int_0^1 z(s)k(t,s)z(t)dsdt + \int_0^1 z(t)^2 dt + 2 \int_0^1 f(t)z(t)dt, \quad (1)$$

where $k(t,s)$ is a symmetric, positive definite kernel, $f(t)$ is a given function, and λ is a sufficiently small nonnegative scalar parameter. The standard variational approach to this problem results in a linear Fredholm integral equation which the optimal function must satisfy.

PROPOSITION 1: The function $u(t)$ which minimizes $W[u, \lambda]$ must satisfy the linear Fredholm integral equation

$$u(t) + f(t) + \lambda \int_0^1 k(t,s)u(s)ds = 0, \quad 0 \leq t \leq 1. \quad (2)$$

PROOF: Denote the optimal function $u(t)$. The arbitrary admissible functions may be written as

$$w(t) = u(t) + \epsilon n(t),$$

where ϵ is an arbitrary scalar, and $n(t)$ is an arbitrary function.

The value of the functional for $w(t)$ may be expressed as

$$W[w; \lambda] = W[u, \lambda] + \epsilon C[u, n; \lambda] + \epsilon^2 D[n; \lambda],$$

where

$$C[u, n; \lambda] = 2 \lambda \int_0^1 \int_0^1 u(s)k(t,s)n(t)dsdt + 2 \int_0^1 u(t)n(t)dt + 2 \int_0^1 f(t)n(t)dt$$

$$D[n, \lambda] = \lambda \int_0^1 \int_0^1 n(s)k(t,s)n(t)dsdt + \int_0^1 n(t)^2 dt.$$

For ϵ sufficiently small the term in ϵ^2 may be ignored. If u is in fact the minimizing solution, then for all ϵ

$$C[u, \eta; \lambda] = 0. \quad (3)$$

Applying the fundamental lemma of the calculus of variations

$$\left(\int_0^1 g(t)h(t)dt = 0 \text{ for arbitrary } h(t) \text{ implies } g(t) = 0 \text{ for all } \right.$$

$0 \leq t \leq 1$ (Ref. 1)) to Eq. (3), since $\eta(t)$ is arbitrary, we get

Eq. (2).

Q.E.D.

Another approach to the minimization problem is to ask how the optimal solution changes as λ varies. This is referred to as parametric imbedding. The standard technique would begin with the linear Fredholm integral equation and convert it into a system of initial valued integro-differential equations. The basic result given in Ref. 2 is

PROPOSITION 2: The function $u(t, \lambda)$ which satisfies the linear Fredholm integral Equation (2) is the solution $u(t, \lambda)$ to the following initial valued integro-differential equations in u , augmented by a resolvent kernel $K(t, s, \lambda)$, and conversely:

$$0 = u_\lambda(t, \lambda) + \psi(t, \lambda) + \int_0^1 K(t, s, \lambda)\psi(s, \lambda)ds \quad (4)$$

$$K_\lambda(t, s, \lambda) = \int_0^1 K(t, s', \lambda)K(s', s, \lambda)ds' \quad (5)$$

$$u(t, 0) = -f(t) \quad (6)$$

$$K(t, s, 0) = -k(t, s) \quad (7)$$

$$\psi(t, s) = \int_0^1 k(t, s)u(s, \lambda)d\lambda \quad (8)$$

$$0 \leq t \leq 1, \quad 0 \leq s \leq 1. \quad (9)$$

The Cauchy system (4)-(9) has proved to be very useful for the computation of the solution of Fredholm integral equations, as well as for the study of the sensitivity of the solution to changes in the parameter λ .

We will now show that the Cauchy system (4)-(9) may be derived directly from the original minimization problem, without ever writing down the Fredholm integral Equation (2). In fact, the solution of the minimization problem could have proceeded even if historically the Fredholm integral equation had never been discovered. The Cauchy system is perfectly adequate for describing the optimal function $u(t)$.

PROPOSITION 3: The function $u(t, \lambda)$ which minimizes $W[u; \lambda]$ satisfies the Cauchy system (4)-(9).

PROOF: Suppose $u(t, \lambda)$ is the minimizing function for parameter value λ , and suppose $u(t, \lambda + d\lambda)$ is the minimizing function for parameter value $\lambda + d\lambda$; then the first variations must satisfy the following inequalities for arbitrary $\epsilon, \eta, \bar{\epsilon}, \bar{\eta}$:

$$\epsilon C[u(t, \lambda), \eta; \lambda] \geq 0 \quad (10)$$

$$\bar{\epsilon} C[u(t, \lambda + d\lambda), \bar{\eta}; \lambda + d\lambda] \geq 0. \quad (11)$$

Since $\epsilon, \eta, \bar{\epsilon}, \bar{\eta}$ are all arbitrary, pick $\bar{\eta} = \eta$ and $\epsilon = -\bar{\epsilon}$ so that (10) and (11) are

$$-\bar{\epsilon} C[u(t, \lambda), \bar{\eta}; \lambda] \geq 0$$

$$\bar{\epsilon} C[u(t, \lambda + d\lambda), \bar{\eta}; \lambda + d\lambda] \geq 0.$$

Add these two inequalities to get

$$\begin{aligned} \bar{\epsilon} [2\lambda \int_0^1 \int_0^1 (u(s, \lambda+d\lambda) - u(s, \lambda)) k(t, s) \bar{\eta}(t) ds dt + 2 \int_0^1 (u(t, \lambda+d\lambda) - u(t, \lambda)) \bar{\eta}(t) dt \\ + 2 d\lambda \int_0^1 \int_0^1 u(s, \lambda+d\lambda) k(t, s) \bar{\eta}(t) ds dt] \geq 0. \end{aligned} \quad (12)$$

Since $\bar{\epsilon}$ has arbitrary sign, the term in square brackets in (12) must be zero. Since $\bar{\eta}(t)$ is arbitrary, apply the fundamental lemma of the calculus of variations to the bracketed term to get

$$\begin{aligned} 2\lambda \int_0^1 k(t, s) (u(s, \lambda+d\lambda) - u(s, \lambda)) ds + 2(u(t, \lambda+d\lambda) - u(t, \lambda)) \\ + 2d\lambda \int_0^1 u(s, \lambda+d\lambda) k(t, s) ds = 0. \end{aligned} \quad (13)$$

Divide (13) by $2d\lambda$ and take the limit as $d\lambda \rightarrow 0$,

$$0 = u_\lambda(t, \lambda) + \lambda \int_0^1 k(t, s) u_\lambda(s, \lambda) ds + \int_0^1 k(t, s) u(s, \lambda) ds. \quad (14)$$

This is a Fredholm integral equation with kernel $k(t, s)$. Corresponding to $k(t, s)$ is a resolvent kernel $K(t, s, \lambda)$ such that solutions of Fredholm integral equations

$$0 = v(t) + \psi(t, \lambda) + \lambda \int_0^1 k(t, s) v(s) ds \quad (15)$$

can be expressed as

$$0 = v(t) + \psi(t, \lambda) + \lambda \int_0^1 K(t, s, \lambda) \psi(s, \lambda) ds.$$

Using the resolvent kernel to express the solution of (14), where

$$\psi(t, \lambda) = \int_0^1 k(t, s) u(s, \lambda) ds, \text{ we get}$$

$$0 = u_\lambda(t, \lambda) + \psi(t, \lambda) + \lambda \int_0^1 K(t, s, \lambda) \psi(s, \lambda) ds.$$

This is nothing other than Eq. (4) of the Cauchy system. The resolvent kernel must itself satisfy a Fredholm integral equation

$$0 = K(t, s, \lambda) + k(t, s) + \lambda \int_0^1 k(t, s') K(s', s, \lambda) ds'. \quad (16)$$

Differentiate (16) with respect to λ to get

$$0 = K_\lambda(t, s, \lambda) + \int_0^1 k(t, s') K(s', s, \lambda) ds' + \lambda \int_0^1 k(t, s') K_\lambda(s', s, \lambda) ds. \quad (17)$$

From Eq. (16) we can reexpress the second term of (17):

$$0 = K_\lambda(t, s, \lambda) - \frac{1}{\lambda} (K(t, s, \lambda) + k(t, s)) + \lambda \int_0^1 k(t, s') K_\lambda(s', s, \lambda) ds'. \quad (18)$$

However, Eq. (18) is a Fredholm integral equation with the same kernel as Eq. (15). Its solution may be expressed using the same resolvent kernel $K(t, s, \lambda)$,

$$0 = K_\lambda(t, s, \lambda) - \frac{1}{\lambda} (K(t, s, \lambda) - k(t, s)) + \lambda \int_0^1 K(t, s, \lambda) - \left(\frac{1}{\lambda}\right) (K(s', s, \lambda) + k(s', s)) ds'. \quad (19)$$

Using Eq. (16) again, several terms in Eq. (19) cancel out, leaving

$$K_\lambda(t, s, \lambda) = \int_0^1 K(t, s', \lambda) K(s', s, \lambda) ds'$$

which is just Eq. (5) of the Cauchy system. To get the initial conditions (7), set $\lambda = 0$ in Eq. (16). To get initial condition (6)

set $\lambda = 0$ in $W[z, \lambda]$. The minimizing function $u(t, 0)$ of

$$\int_0^1 u(t, 0)^2 dt + 2 \int_0^1 f(t) u(t, 0) dt$$

can easily be shown to be $u(t, 0) = f(t)$.

Q.E.D.

3. DISCUSSION

The main purpose of this paper has been to reduce the quadratic variational problem in Eq. (1) to the Cauchy system in relations (4)-(9). In particular we have been able to do this without making any use of the Euler equation, which takes the form of the Fredholm integral equation (2).

Subsequent papers in this series will be devoted to systems and to the treatment of non-quadratic variational problems. Application to team decision theory and other areas will be presented.

REFERENCES

1. COURANT, R. and HILBERT, D., Methods of Mathematical Physics, Interscience, New York, 1953.
2. KAGIWADA, H. and KALABA, R., Integral Equations via Imbedding Methods, Addison-Wesley Publishers, Inc., Reading, Massachusetts, 1974.
3. CASTI, J. and KALABA, R., Imbedding Methods in Applied Mathematics, Addison-Wesley Publishers, Inc., Reading, Massachusetts, 1973.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		REAL INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR-78-0016	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) DIRECT TRANSFORMATION OF VARIATIONAL PROBLEM INTO CAUCHY SYSTEMS. I. SCALAR-QUADRATIC CASE.		5. TYPE OF REPORT & PERIOD COVERED Interim rept.
7. AUTHOR(s) James/Hess Robert/Kalaba		6. PERFORMING ORG. REPORT NUMBER MRG-7710
		8. CONTRACT OR GRANT NUMBER(s) AFOSR-77-3383
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Southern California Department of Economics Los Angeles, CA 90007		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102 2304 A3
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Rsch/NM Bolling AFB, DC 20332		12. REPORT DATE 11 1977
		13. NUMBER OF PAGES 12 13 p.
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) USCMRG-7710		15. SECURITY CLASS. (of this report) UNCLASSIFIED
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) variational problems, integral equations, parametric imbedding		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This series of papers addresses three interrelated problems: the solution of a variational minimization problem, the solution of integral equations, and the solution of an initial valued system of integrodifferential equations. It will be shown that a large class of minimization problems requires the solution of linear Fredholm integral equations. It has also been shown that the solution of a linear Fredholm integral equation is identical to the solution of a Cauchy system. In this paper, we by-pass the Fredholm integral equations and show that the minimization problem directly implies a solution of a Cauchy system. This		

20. Abstract

↓
· (continued) first paper in the series looks only at quadratic functionals
and scalar functions. ↑

UNCLASSIFIED